

Enhancement of parametric pumping due to Andreev reflection

Jian Wang¹, Yadong Wei¹, Baigeng Wang¹, Hong Guo²

1. Department of Physics, The University of Hong Kong, Pokfulam Road, Hong Kong, China

2. Center for the Physics of Materials and Department of Physics, McGill University, Montreal, PQ, Canada H3A 2T8

Abstract

We report properties of parametric electron pumping in the presence of a superconducting lead. Due to a constructive interference between the direct reflection and the multiple Andreev reflection, the pumped current is greatly enhanced. For both quantum point contacts and double barrier structures at resonance, we obtain exact solutions in the weak pumping regime showing that $I_p^{NS} = 4I_p^N$, which should be compared with the result of conductance $G_{NS} = 2G_N$. Numerical results are also provided for the strong pumping regime showing interesting Andreev assisted pumping behaviour.

73.23.Ad,73.40.Gk,72.10.Bg,74.50.+r

Current can flow under zero bias when two system parameters of a nanostructure are varied in a cyclic fashion. The physics of this parametric electron pump has been analyzed by several authors [1,2]. Recently, it has been realized experimentally by Switkes *et. al.* [3] confirming many of the theoretical predictions. So far, investigations on parametric pumping are focused on normal nanostructures. It will be interesting to study a hybrid structure where a superconducting lead is present. In the presence of normal conductor-superconductor (NS) interface, an incoming electron-like excitation can be Andreev reflected as a hole-like excitation. This current doubling effect gives rise to the relation for conductance $G_{NS} = 2G_N$ for hybrid quantum point contacts and quantum dots at resonance [4]. For dirty NS contacts, de Jong and Beenakker [5] found that the shot noise at subgap voltages is also doubled with respect to its value in normal state which has been confirmed experimentally [6]. To further explore how does Andreev reflection modify quantum interference in the normal state, we have investigated the parametric pumping phenomenon in the presence of a superconducting lead. We find that due to quantum interference of the direct reflection and the multiple Andreev reflection, the pumped current is greatly enhanced. For quantum point contact and quantum dot at resonance, we obtained a relation for the pumped current, $I_p^{NS} = 4I_p^N$ in the weak pumping regime. Numerical results are presented in the strong pumping regime showing interesting Andreev assisted pumping behaviour which can be verified experimentally.

We consider a parametric pump which consists of a double barrier tunneling structure attached to a normal left lead and a superconducting right lead. The double barrier structure is modeled by potential $V(x) = V_1\delta(x+a) + V_2\delta(x-a)$ where $V_1 = V_0 + V_p \sin(\omega t)$ and $V_2 = V_0 + V_p \sin(\omega t + \phi)$ and V_p is the pumping amplitude. We further apply a gate voltage v_g to control the energy level of the structure. The units are fixed by setting $\hbar = 2m = q = 1$ in the following analysis [7]. At low frequencies, the adiabatic pumped current in the presence of superconducting leads is [1,8]

$$I_p^{NS} = \frac{\omega}{\pi} \int_0^\tau d\tau \left[\frac{dN_L}{dV_1} \frac{dV_1}{dt} + \frac{dN_L}{dV_2} \frac{dV_2}{dt} \right] \quad (1)$$

where the quantity dN_L/dV is the injectivity [9,10] given, at zero temperature, by

$$\frac{dN_L}{dV_j} = \frac{1}{2\pi} \text{Im}[S_{ee}^* \partial_{V_j} S_{ee} - S_{he}^* \partial_{V_j} S_{he}] \quad (2)$$

where the first term is the injectivity of electron due to the external parameter [9,10], *i.e.* the partial density of states (DOS) for an electron coming from left lead and exiting the system as an electron, and the second term is the injectivity of a hole, *i.e.* the DOS for a hole coming from left lead and exiting the system as an electron.

For the hybrid nanostructure, the scattering matrix S_{ee} and S_{he} are given by [4,11]

$$\hat{S} = \hat{S}_{11} + \hat{S}_{12}(1 - \hat{R}_I \hat{S}_{22})^{-1} \hat{R}_I \hat{S}_{21} \quad (3)$$

where $\hat{S}$ is a 2×2 scattering matrix for NS structure with matrix element $S_{\mu\nu}$ ($\mu, \nu = e, h$) and $\hat{S}_{ij}(E)$ ($i, j = 1, 2$) is a diagonal 2×2 scattering matrix for double barrier structure with matrix element $S_{ij}(E)$ and $S_{ij}^*(-E)$. $\hat{R}_I = \alpha \sigma_x$ is the 2×2 scattering matrix at NS interface with off diagonal matrix element α . Here $\alpha = (E - i\nu\sqrt{\Delta^2 - E^2})/\Delta$ with $\nu = 1$ when $E > -\Delta$ and $\nu = -1$ when $E < -\Delta$. In Eq.(3), the energy E is measured relative to the chemical potential μ of the superconducting lead. Eq.(3) has clear physical meaning [11]. The first term is the direct reflection from the normal scattering structure and the second term can be expanded as $\hat{S}_{12}\hat{R}_I\hat{S}_{21} + \hat{S}_{12}\hat{R}_I\hat{S}_{22}\hat{R}_I\hat{S}_{21} + \dots$ which is clearly the multiple Andreev reflection in the hybrid structure. From Eq.(3) we obtain the well known expressions for the scattering matrix S_{ee} and S_{he} [4]

$$S_{ee}(E) = S_{11}(E) + \alpha^2 S_{12}(E) S_{22}^*(-E) M_e S_{21}(E) \quad (4)$$

and

$$S_{he}(E) = \alpha S_{12}^*(-E) M_e S_{21}(E) \quad (5)$$

with $M_e = [1 - \alpha^2 S_{22}(E) S_{22}^*(-E)]^{-1}$. In the following, we first present the exact result for the pumped current in the weak pumping regime and then study the general situation numerically. In the weak pumping regime, Eq.(1) can be expanded to the lowest order in V_p ,

$$I_p^{NS} = \frac{\omega \sin \phi V_p^2}{\pi} \text{Im}[\partial_{V_1} S_{ee}^* \partial_{V_2} S_{ee} - \partial_{V_1} S_{he}^* \partial_{V_2} S_{he}] \quad (6)$$

as compared with the expression for the normal structure [1,12],

$$I_p^N = \frac{\omega \sin \phi V_p^2}{\pi} \text{Im}[\partial_{V_1} S_{11}^* \partial_{V_2} S_{11} + \partial_{V_1} S_{21}^* \partial_{V_2} S_{21}] \quad (7)$$

where we set $V_p = 0$ in $S_{\nu\mu}$ and S_{ij} after the partial derivatives. We further assume that the Fermi energy is close to the chemical potential of superconducting lead, so $E \sim 0$ and $\alpha \sim -i$. Under this condition, S_{he} and hence $\partial_{V_j} S_{he}$ are pure imaginary numbers for general $V_j(t)$. As a result, S_{he} does not contribute to I_p^{NS} in Eq.(1) as long as $E = 0$ [13]. The only contribution comes from S_{ee} which is the superposition of the direct reflection and the multiple Andreev reflection. We will consider two cases: (a) a quantum point contact, *e.g.* $V_0 = 0$; and (b) the double barrier quantum dot at resonance. For both cases, $S_{11} = 0$ in the absence of pumping voltage. Therefore, from Eqs.(4) we have

$$\partial_{V_{1/2}} S_{ee} = \partial_{V_{1/2}} S_{11} - S_{12}^2 \partial_{V_{2/1}} S_{11}^* \quad (8)$$

where we have used the fact that $\partial_{V_1} S_{22} = \partial_{V_2} S_{11}$. Using Fisher-Lee relation [14] $S_{\alpha\beta} = -\delta_{\alpha\beta} + i\sqrt{v_\alpha v_\beta} G_{\alpha\beta}^r$ and the Dyson equation $\partial_{V_j} G_{\alpha\beta}^r = G_{\alpha j}^r G_{j\beta}^r$ [15], we have $\partial_{V_1} S_{11} = iv G_{11}^r G_{11}^r = -i/(2k)$ and $\partial_{V_2} S_{11} = iv G_{12}^r G_{21}^r = -i/(2k) S_{12}^2$ with the velocity $v = 2k$. So for both cases (a) and (b) we have constructive interference between direct reflection and the multiple Andreev reflection $\partial_{V_j} S_{ee} = 2\partial_{V_j} S_{11}$ which gives a pumped current $I_p^{NS} = 4I_p^N$ with $I_p^N = \pm\omega V_p^2 \sin(4ka)/(4\pi k^2)$, where the plus sign is for quantum point contact since $S_{12} = e^{2ika}$, and the minus sign corresponds to resonant tunneling since $S_{12} = e^{-2ika}$.

In the general situation, the pumped current can be calculated numerically using Eq.(1) [16]. Since the pumped current is proportional to ω , we set $\omega = 1$ for convenience. In the left inset of Fig.1, we plot the ratio I_p^{NS}/I_p^N as a function of pumping strength V_p/V_0 for $V_0 = 79.2$ at the resonant point. We see that as the pumping amplitude increases, the constructive interference effect is suppressed. At small pumping strength, the ratio is about four which

agrees with our theoretical analysis. At larger pumping strength, this ratio decreases to the value below two. Similar behaviour is seen for the quantum point contact. Fig.1 shows the pumped current as a function of Fermi energy E_F for different pumping amplitudes. For $V_0 = 79.2$, we have chosen $v_g = -9.39$ so that one resonant level in the quantum dot is aligned with the chemical potential μ_s of the superconducting lead in the absence of pumping voltage V_p . For comparison, we also plot the Andreev reflection coefficient when $V_p = 0$. Several observations are in order: (1). the pumped current is peaked near the resonant level showing clearly a resonant behaviour. This is because the pumped current (Eq.(1) is proportional to DOS of the system which reaches its maximum near the resonance. (2) As the pumping amplitude V_p increases, the pumped current increases. (3). The pumped current has two asymmetric peaks. To understand this, we plot in the right inset of Fig.1 the Andreev reflection coefficient versus E_F at different moments in one pumping cycle. From this inset, we observe that the Andreev reflection coefficient gives one or two peaks depending on the configuration of the system. This behaviour can be understood from the Breit-Wigner form of the resonant Andreev reflection T_A through a single level $E_0 = 0$ (measured relative to μ_s) [17]:

$$T_A = \frac{\Gamma_L^2 \Gamma_R^2}{4(E^2 + \Gamma \Delta_\Gamma/4)^2 + \Gamma_L^2 \Gamma_R^2} \quad (9)$$

where $\Delta_\Gamma = \Gamma_L - \Gamma_R$ and $\Gamma = \Gamma_L + \Gamma_R$. We see that T_A shows two peaks when $\Gamma_L < \Gamma_R$ and just one peak otherwise. Note that in one pumping cycle half of the configurations corresponds to $\Gamma_L < \Gamma_R$, therefore two pumped current peaks show up in Fig.1 because from Eq.(1) the pumped current is obtained through integral over all the configurations in one pumping cycle. Finally, the reason that two peaks are asymmetric is mainly due to the energy dependence of the self energy. If μ_s is right in the middle of two resonant levels (E_1 and E_2), *i.e.*, $\mu = (E_1 + E_2)/2$, then electron coming from normal lead with incident energy E_1 tunnels into the structure through the resonant level E_1 and Andreev reflected as a hole back to the quantum dot through the resonant level E_2 with a Cooper pair created in the superconducting lead. We now examine the Andreev assisted pumping through two

levels. For $V_0 = 79.2$ and $v_g = -23.48$, there are two resonant levels inside the subgap ($\mu_s = 0$) at $E_1 = 14.09$ and $E_2 = -14.09$. Hence, strong Andreev reflections can occur near $E_F = E_1$. Fig.2 shows the two level pumped current versus Fermi energy (solid line). For comparison we also plot the corresponding Andreev reflection coefficient versus Fermi energy when the V_p is switched off (see inset of Fig.2). Similar to Fig.1, the pumped current also shows strong resonant behaviour with smaller amplitude (compare Fig.1 dot-dashed line). We found two peaks of pumped current around E_1 , one is near the resonant energy and the other one is shifted to a smaller energy with larger current. Although it is similar to Fig.1 but has different origin. This is due to the fact that when the pumping gate is turned on, the barrier heights and hence the resonant level change with time. To confirm this, we also plot the Andreev reflection coefficients T_A at several instants of one pumping cycle in the same figure. The peaks of T_A shift around the energy level at different pumping time and give the behaviour of pumped current. We have also calculated the pumped current for other system parameters and confirmed that the behaviour of pumped current shown here is generic.

In summary, in the presence of superconducting lead, the pumped current is greatly enhanced due to the quantum interference of direct reflection and multiple Andreev reflection. In the weak pumping regime, we have the relation $I_p^{NS} = 4I_p^N$ for both quantum point contact and the resonant tunneling structure. Interesting Andreev assisted pumping behaviours are revealed as well.

We gratefully acknowledge support by a RGC grant from the SAR Government of Hong Kong under grant number HKU 7215/99P. H.G. is supported by NSERC of Canada and FCAR of Québec.

REFERENCES

- [1] P.W. Brouwer, Phys. Rev. B **58**, R10135 (1998).
- [2] F. Zhou, B. Spivak, and B.L. Altshuler, Phys. Rev. Lett. **82**, 608 (1999). T.A. Shutenko, I.L. Aleiner, and B.L. Altshuler, Phys. Rev. B. **61**, 10366 (2000).
- [3] M. Switkes, C. Marcus, K. Capman, and A.C. Gossard, Science **283**, 1905 (1999).
- [4] C.W.J. Beenakker, Rev. Mod. Phys. **69**, 731 (1997).
- [5] M.J.M. de Jong and C.W.J. Beenakker, Phys. Rev. B **49**, 16070 (1994).
- [6] X. Jehl, P. Payet-Burin, C. Baraduc, R. Calemczuk, and M. Sanquer, Phys. Rev. Lett. **83**, 1660 (1999). X. Jehl, M. Sanquer, R. Calemczuk, and D. Mailly, Nature (London) **405**, 50 (2000).
- [7] For the GaAs system with $a = 1000A$, the energy unit is $E = 56\mu eV$.
- [8] This formula can be derived using the Keldysh nonequilibrium Green's function method used in the present work.
- [9] T. Gramespacher and M. Buttiker, Phys. Rev. B **61**, 8125 (2000).
- [10] J. Wang, Y.D. Wei, H. Guo, Q.F. Sun, and T.H. Lin, cond-mat/0011150.
- [11] G.B. Lesovik, A.L. Fauchere, and G. Blatter, Phys. Rev. B **55**, 3146 (1997).
- [12] A factor of two has been included for spin.
- [13] Here we assume a zero magnetic field. Since S_{12} is symmetric with respect to V_1 and V_2 , it does not contribute to I_p^N .
- [14] D. S. Fisher and P. A. Lee, Phys. Rev. B **23**, 6851 (1981).
- [15] V. Gasparian, T. Christen, and M. Buttiker, Phys. Rev. A **54**, 4022 (1996).
- [16] For the double barrier structure, the scattering matrices can be easily obtained. See

for example, M.K. Yip, J. Wang, and H. Guo, Z. Phys. B: Condens. Matter **104**, 463 (1997).

[17] Y.D. Wei, J. Wang, H. Guo, H. Mehrez, and C. Roland, cond-mat/0103210, to appear in Phys. Rev. B. (May 15 issue, 2001).

[18] Y.D. Wei, J. Wang, and H. Guo, Phys. Rev. B **62**, 9947 (2000).

FIGURES

FIG. 1. The pumped current versus E_F at different pumping amplitudes: $V_p = 0.03V_0$, $0.05V_0$. The left inset: I_p^{NS}/I_p^N versus V_p/V_0 at resonant point. The right inset: Andreev reflection coefficient T_A as a function of E_F with $V_p = 0.05V_0$ at different pumping time: $t = \pi/4$ (dot dashed line), $t = \pi/2$ (dotted line), $t = 3\pi/4$ (solid line), and $t = 7\pi/4$ (dashed line). System parameters in Fig.1 and the insets: $V_0 = 79.2$, $v_g = -9.39$, $\phi = \pi/2$, and $\Delta = 20$.

FIG. 2. The two level pumped current I_p as a function of Fermi energy (solid line) with $V_p = 0.05V_0$ and $v_g = -23.481$. We also plot T_A at several pumping time: $t = \pi/4$ (dotted line), $t = \pi/2$ (dashed line), $t = 3\pi/4$ (dot-dashed line), and $t = \pi$ (long dashed line). The pumped current has been offset by 0.3 for illustrating purpose. Inset: T_A versus Fermi energy when $V_p = 0$. Other parameters are the same as Fig.1.

Fig1



